

AD704283

FTD-HT-23-635-69


FOREIGN TECHNOLOGY DIVISION



USE OF BLOOD AGAR WITH SALT FOR SEPARATING TOXIGENIC
STAPHYLOCOCCI FROM AIR

by

L. S. Sarochinskaya



D D C
REF ID: A64111
APR 24 1970
RUSSELL C

Distribution of this document is unlimited. It may be released to the Clearinghouse, Department of Commerce, for sale to the general public.

Reproduced by the
CLEARINGHOUSE
for Federal Scientific & Technical
Information Springfield Va. 22151

EDITED TRANSLATION

USE OF BLOOD AGAR WITH SALT FOR SEPARATING TOXIGENIC STAPHYLOCOCCI FROM AIR

By: L. S. Sarochinskaya

English pages: 2

Source: Laboratornoye Delo (Laboratory Work), No. 3, 1964, p. 167.

Translated by: L. Thompson/TDBRS-3

THIS TRANSLATION IS A REPRODUCTION OF THE ORIGINAL FOREIGN TEXT WITHOUT ANY ANALYTICAL OR EDITORIAL COMMENT. STATEMENTS OR THEORIES ADVOCATED OR IMPLIED ARE THOSE OF THE SOURCE AND DO NOT NECESSARILY REFLECT THE POSITION OR OPINION OF THE FOREIGN TECHNOLOGY DIVISION.

PREPARED BY:

TRANSLATION DIVISION
FOREIGN TECHNOLOGY DIVISION
WP-APB, OND.

USE OF BLOOD AGAR WITH SALT FOR SEPARATING TOXIGENIC
STAPHYLOCOCCI FROM AIR

L. S. Sarochinskaya

Department of General Hygiene I, I. P. Pavlov
Medical Institute, Leningrad

In evaluating the microbial seeding of the air in closed areas, it is necessary to study the total number of microorganisms, the number of Streptococci viridans, Streptococci hemolyticus, and toxigenic Staphylococci per m^3 of air¹.

For detection of toxigenic Staphylococci in the air of closed areas, it is recommended that a sample of air be taken on lactic-salt agar, and the toxigenic properties of the colony of Staphylococci growing on that medium be checked, studying their hemolytic capacity and a group of other signs of toxigenicity.

In tests of the air of hospital areas, up to 18-26 colonies of Staphylococci on one dish grew on the dishes of the primary seeding. For this reason, 18-26 dishes with blood agar would be necessary for studying the hemolytic capacity of Staphylococci growing on a single primary seeding dish. In practice this is not feasible, and consequently a quantitative estimate of toxigenic Staphylococci in the air by the given method is not possible.

¹Procedural instructions on sanitary-bacteriological investigation of external environment objects. Moscow, 1961.

Research is significantly facilitated, its time is reduced, the nutrient medium is conserved, and more reliable results are obtained if 5% blood agar with 5-6.5% salt is used for detecting Staphylococci in the air.

Before the test, 2.5% meat-peptone agar with 5-6.5% salt (pH 7.6-7.8) is prepared, and blood is added the moment the preparation is poured into the dishes. It is better to use rabbit blood, which is more sensitive¹, but good results are also obtained with human placental and citrate blood.

The growth picture of Staphylococci on this medium is so typical that even a researcher with little experience in work, encounters no difficulties in processing the cultures.

Thus, by taking samples of air on blood agar with salt, we have the opportunity to calculate the amount of toxigenic Staphylococci per unit volume of air as early as the end of the second day, and then to continue the furthest study of toxigenicity of the cultures of Staphylococci which are formed.

In using blood agar with salt for separating toxigenic Staphylococci from air, we checked more than a thousand studies of hospital-area air and were satisfied of the advantages of this medium for initial seedings of air as compared to other culture mediums.

Submitted
24 December 1962

¹K. I. Turzhetskii, Doctoral dissertation, Leningrad, 1950.

DATA HANDLING PAGE				
01-ACCESSION NO. TT0000222		02-DOCUMENT LOC		
03-TITLE USE OF BLOOD AGAR WITH SALT FOR SEPARATING TOXIGENIC STAPHYLOCOCCI FROM AIR -II-		04-TOPIC TAGS medical R and D, atmospheric contamination, oxygen toxicity, staphylococcus, blood agent		
05-SUBJECT AREA 06				
07-AUTHOR CO-AUTHORS SAROCHINSKAYA, L. S.				08-DATE OF INFO -----64
09-SOURCE LABORATORNOYE DELO, MOSCOW, MEDITSINA (RUSSIAN)				10-DOCUMENT NO. FTD-HT-23-635-69
				11-PROJECT NO. 6030204
12-SECURITY AND DOWNGRADING INFORMATION UNCL. 0			13-CONTROL MARKINGS NONE	14-HEADER CLASS UNCL
15-REEL/FRAME NO. 1891 0814	16-SUPERSEDES	17-CHANGES	18-GEOGRAPHICAL AREA UR	19-NO. OF PAGES 2
20-CONTRACT NO.	21-X REF ACC. NO. 45-	22-PUBLISHING DATE 94-	23-TYPE PRODUCT TRANSLATION	24-REVISION FREQ NONE
25-STEP NO. 02-UR/0000/64/000/003/0167/0167				
26-ABSTRACT (U) The article describes the use of blood agar with 5-6.5 percent salt (pH 7.6-7.8) for calculating the amount of toxigenic staphylococci per unit volume of air in closed areas. The advantages of using this medium include: reduced time, conservation of the medium, and more reliable results as compared to other culture mediums.				